


TReitement des Images pour la Vision Artificielle

Vincent Lepetit

slides in part based on material from Mathieu Aubry, Andrew Zisserman, David Lowe

Bayesian Theory for Computer Vision

In the previous lecture



$$\arg \min_{\mathbf{R}, \mathbf{T}} \frac{1}{|\mathcal{M}|} \sum_i \|\text{proj}(\mathbf{M}_i; \mathbf{R}, \mathbf{T}) - \mathbf{m}_i\|^2$$

In this lecture

$$\arg \max_{\mathbf{x}} p(\mathbf{x} \mid \mathbf{z})$$

\mathbf{x} : state [in our case, for example \mathbf{R} , \mathbf{T}]

\mathbf{z} : observations [in our case, for example feature points \mathbf{m}_i]

$p(\mathbf{x} \mid \mathbf{z})$: probability density function of \mathbf{x} given \mathbf{z}

probability density function

$p(\mathbf{x})$: probability density function (pdf) of \mathbf{x}

Intuitively:


- $p(\mathbf{x}) \sim$ how likely the random variable for \mathbf{x} is close to \mathbf{x}
- $p(\mathbf{x}_1) > p(\mathbf{x}_2)$: the correct value is more likely to be \mathbf{x}_1 than \mathbf{x}_2

probability density function (2)

For a univariate random variable X :

$$\Pr(a \leq X \leq b) = \int_a^b p(x)dx$$


$$p(x) = N(x; 0, 0.2)$$



$$\Pr(a \leq X \leq b) = \int_a^b p(x)dx$$

probability density function (2)

For a 2D random variable:



$$\arg \max_{\mathbf{x}} p(\mathbf{x} | \mathbf{z})$$

$$p(\mathbf{x} | \mathbf{z})$$

Bayes' theorem

$$\arg \max_{\mathbf{x}} p(\mathbf{x} \mid \mathbf{z})$$

\mathbf{x} : state [in our case, for example \mathbf{R} , \mathbf{T}]

\mathbf{z} : observations [in our case, for example feature points \mathbf{m}_i]

$p(\mathbf{x} \mid \mathbf{z})$: probability density function of \mathbf{x} given \mathbf{z}

$$p(\mathbf{x} \mid \mathbf{z}) = \frac{1}{p(\mathbf{z})} p(\mathbf{z} \mid \mathbf{x}) p(\mathbf{x})$$

posterior likelihood prior

Likelihood - example


$$p(\mathbf{z} \mid \mathbf{x}) = p(\mathbf{z}_1, \dots, \mathbf{z}_n \mid \mathbf{x})$$

$$= \prod_i p(\mathbf{z}_i \mid \mathbf{x})$$


if we assume
the observations \mathbf{z}_i
are independent given \mathbf{x}

$$= \prod_i N(\mathbf{m}_i; \text{proj}(\mathbf{M}_i; \mathbf{R}, \mathbf{T}), \Sigma)$$

if we assume the noise
on the observations
is Gaussian, with covariance Σ



Prior



If we have no idea what the state can be:

$$p(\mathbf{x}) = \mu$$

If we believe the camera center is close to Point $(0,0,0)$:

$$p(\mathbf{x}) = N(\mathbf{C}(\mathbf{x}); \mathbf{0}, \Sigma_{\mathbf{x}})$$

etc.

Link (1)

$$\begin{aligned} & \arg \max_{\mathbf{x}} p(\mathbf{x} \mid \mathbf{z}) \\ = & \arg \max_{\mathbf{x}} \frac{1}{p(\mathbf{z})} p(\mathbf{z} \mid \mathbf{x}) p(\mathbf{x}) \\ = & \arg \max_{\mathbf{R}, \mathbf{T}} \prod_i N(\mathbf{m}_i; \text{proj}(\mathbf{M}_i; \mathbf{R}, \mathbf{T}), \Sigma) \\ = & \arg \min_{\mathbf{R}, \mathbf{T}} -\log \prod_i N(\mathbf{m}_i; \text{proj}(\mathbf{M}_i; \mathbf{R}, \mathbf{T}), \Sigma) \\ = & \arg \min_{\mathbf{R}, \mathbf{T}} -\sum_i \log N(\mathbf{m}_i; \text{proj}(\mathbf{M}_i; \mathbf{R}, \mathbf{T}), \Sigma) \\ = & \arg \min_{\mathbf{R}, \mathbf{T}} \sum_i \|\mathbf{m}_i - \text{proj}(\mathbf{M}_i; \mathbf{R}, \mathbf{T})\|^2 \end{aligned}$$

Link (2)

$$\begin{aligned} & \arg \max_{\mathbf{x}} p(\mathbf{x} \mid \mathbf{z}) \\ = & \arg \max_{\mathbf{x}} \frac{1}{p(\mathbf{z})} p(\mathbf{z} \mid \mathbf{x}) p(\mathbf{x}) \\ = & \arg \max_{\mathbf{R}, \mathbf{T}} \prod_i N(\mathbf{m}_i; \text{proj}(\mathbf{M}_i; \mathbf{R}, \mathbf{T}), \Sigma) N(\mathbf{C}(\mathbf{R}, \mathbf{T}); \mathbf{0}, \Sigma_{\mathbf{x}}) \\ = & \arg \min_{\mathbf{R}, \mathbf{T}} - \sum_i \log N(\mathbf{m}_i; \text{proj}(\mathbf{M}_i; \mathbf{R}, \mathbf{T}), \Sigma) + \log N(\mathbf{C}(\mathbf{R}, \mathbf{T}); \mathbf{0}, \Sigma_{\mathbf{x}}) \\ = & \arg \min_{\mathbf{R}, \mathbf{T}} \underbrace{\sum_i \|\mathbf{m}_i - \text{proj}(\mathbf{M}_i; \mathbf{R}, \mathbf{T})\|^2}_{\text{log-likelihood}} + \underbrace{\lambda \|\mathbf{C}(\mathbf{R}, \mathbf{T})\|^2}_{\text{regularization term}} \end{aligned}$$

Robust estimators


If we assume the noise on the observations is Gaussian:

$$p(\mathbf{z}_i \mid \mathbf{x}) = N(\mathbf{m}_i; \text{proj}(\mathbf{M}_i; \mathbf{R}, \mathbf{T}), \Sigma)$$


If we consider that \mathbf{m}_i can be either an inlier with Gaussian noise or an outlier, we can take:

$$p(\mathbf{z}_i \mid \mathbf{x}) \sim N(\mathbf{m}_i; \text{proj}(\mathbf{M}_i; \mathbf{R}, \mathbf{T}), \Sigma) + \mu$$

Normal distribution
(inliers)




Uniform distribution
(outliers)




+

=

Mixture




$-\log(\cdot)$




Least-squares

$-\log(\cdot)$



Tukey estimator

Tracking over Time



Tracking over Time

\mathbf{x}_t : State at time t

$\mathbf{X}_t = \mathbf{x}_t, \mathbf{x}_{t-1}, \dots, \mathbf{x}_0$: States from time 0 to time t


\mathbf{z}_t : Observations performed at time t .

$\mathbf{Z}_t = \mathbf{z}_t, \mathbf{z}_{t-1}, \dots, \mathbf{z}_0$: Observations up to time t

How can we estimate the density $p(\mathbf{x}_t | \mathbf{Z}_t)$?

ie What is the density on the object state at time t given the images captured up to time t ?

Example



Tracking over Time

\mathbf{x}_t : State at time t

$\mathbf{X}_t = \mathbf{x}_t, \mathbf{x}_{t-1}, \dots, \mathbf{x}_0$: States from time 0 to time t

\mathbf{z}_t : Observations performed at time t .

$\mathbf{Z}_t = \mathbf{z}_t, \mathbf{z}_{t-1}, \dots, \mathbf{z}_0$: Observations up to time t

How can we estimate the density $p(\mathbf{x}_t | \mathbf{Z}_t)$?

i.e. What is the density on the object state at time t given the images captured up to time t ?

We will see that:

$$p(\mathbf{x}_t | \mathbf{Z}_t) \sim \underbrace{p(\mathbf{z}_t | \mathbf{x}_t)}_{\substack{\text{likelihood /} \\ \text{Observation model}}} \int_{\mathbf{x}_{t-1}} \underbrace{p(\mathbf{x}_t | \mathbf{x}_{t-1})}_{\text{Motion model}} \underbrace{p(\mathbf{x}_{t-1} | \mathbf{Z}_{t-1})}_{\text{Density estimated at time } t-1}$$

Tracking over Time

\mathbf{x}_t : State at time t

$\mathbf{X}_t = \mathbf{x}_t, \mathbf{x}_{t-1}, \dots, \mathbf{x}_0$: States from time 0 to time t

\mathbf{z}_t : Observations performed at time t .

$\mathbf{Z}_t = \mathbf{z}_t, \mathbf{z}_{t-1}, \dots, \mathbf{z}_0$: Observations up to time t

$$p(\mathbf{X}_t, \mathbf{Z}_t) = p(\mathbf{x}_t, \mathbf{z}_t, \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) \quad [\text{definition}]$$

$$= p(\mathbf{x}_t, \mathbf{z}_t \mid \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) \quad [\text{conditional probability: } p(A, B) = p(A \mid B) p(B)]$$

$$= p(\mathbf{z}_t \mid \mathbf{x}_t, \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{x}_t \mid \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) \quad [\text{conditional probability}]$$

Tracking over Time

\mathbf{x}_t : State at time t

$\mathbf{X}_t = \mathbf{x}_t, \mathbf{x}_{t-1}, \dots, \mathbf{x}_0$: States from time 0 to time t

\mathbf{z}_t : Observations performed at time t .

$\mathbf{Z}_t = \mathbf{z}_t, \mathbf{z}_{t-1}, \dots, \mathbf{z}_0$: Observations up to time t

We just saw:

$$p(\mathbf{X}_t, \mathbf{Z}_t) = p(\mathbf{z}_t \mid \mathbf{x}_t, \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{x}_t \mid \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{X}_{t-1}, \mathbf{Z}_{t-1})$$

We also have:

$$p(\mathbf{X}_t, \mathbf{Z}_t) = p(\mathbf{X}_t \mid \mathbf{Z}_t) p(\mathbf{Z}_t)$$

so

$$p(\mathbf{X}_t \mid \mathbf{Z}_t) p(\mathbf{Z}_t) = p(\mathbf{z}_t \mid \mathbf{x}_t, \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{x}_t \mid \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{X}_{t-1}, \mathbf{Z}_{t-1})$$

Tracking over Time

\mathbf{x}_t : State at time t

$\mathbf{X}_t = \mathbf{x}_t, \mathbf{x}_{t-1}, \dots, \mathbf{x}_0$: States from time 0 to time t

\mathbf{z}_t : Observations performed at time t .

$\mathbf{Z}_t = \mathbf{z}_t, \mathbf{z}_{t-1}, \dots, \mathbf{z}_0$: Observations up to time t

$$p(\mathbf{X}_t | \mathbf{Z}_t) p(\mathbf{Z}_t) = p(\mathbf{z}_t | \mathbf{x}_t, \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{x}_t | \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{X}_{t-1}, \mathbf{Z}_{t-1})$$

We also have:

$$p(\mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) = p(\mathbf{X}_{t-1} | \mathbf{Z}_{t-1}) p(\mathbf{Z}_{t-1})$$

so

$$p(\mathbf{X}_t | \mathbf{Z}_t) p(\mathbf{Z}_t) = p(\mathbf{z}_t | \mathbf{x}_t, \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{x}_t | \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{X}_{t-1} | \mathbf{Z}_{t-1}) p(\mathbf{Z}_{t-1})$$

Tracking over Time

\mathbf{x}_t : State at time t

$\mathbf{X}_t = \mathbf{x}_t, \mathbf{x}_{t-1}, \dots, \mathbf{x}_0$: States from time 0 to time t

\mathbf{z}_t : Observations performed at time t .

$\mathbf{Z}_t = \mathbf{z}_t, \mathbf{z}_{t-1}, \dots, \mathbf{z}_0$: Observations up to time t

$$p(\mathbf{X}_t | \mathbf{Z}_t) p(\mathbf{Z}_t) = p(\mathbf{z}_t | \mathbf{x}_t, \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{x}_t | \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{X}_{t-1} | \mathbf{Z}_{t-1}) p(\mathbf{Z}_{t-1})$$

$p(\mathbf{Z}_t)$ is constant with respect to \mathbf{x}_t , so

$$p(\mathbf{X}_t | \mathbf{Z}_t) \sim p(\mathbf{z}_t | \mathbf{x}_t, \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{x}_t | \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{X}_{t-1} | \mathbf{Z}_{t-1})$$

Tracking over Time

\mathbf{x}_t : State at time t

$\mathbf{X}_t = \mathbf{x}_t, \mathbf{x}_{t-1}, \dots, \mathbf{x}_0$: States from time 0 to time t

\mathbf{z}_t : Observations performed at time t .

$\mathbf{Z}_t = \mathbf{z}_t, \mathbf{z}_{t-1}, \dots, \mathbf{z}_0$: Observations up to time t

$$p(\mathbf{X}_t | \mathbf{Z}_t) \sim p(\mathbf{z}_t | \mathbf{x}_t, \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{x}_t | \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{X}_{t-1} | \mathbf{Z}_{t-1})$$

We want to get to:

$$p(\mathbf{x}_t | \mathbf{Z}_t) \sim p(\mathbf{z}_t | \mathbf{x}_t) \int_{\mathbf{x}_{t-1}} p(\mathbf{x}_t | \mathbf{x}_{t-1}) p(\mathbf{x}_{t-1} | \mathbf{Z}_{t-1})$$

Tracking over Time

\mathbf{x}_t : State at time t

$\mathbf{X}_t = \mathbf{x}_t, \mathbf{x}_{t-1}, \dots, \mathbf{x}_0$: States from time 0 to time t

\mathbf{z}_t : Observations performed at time t .

$\mathbf{Z}_t = \mathbf{z}_t, \mathbf{z}_{t-1}, \dots, \mathbf{z}_0$: Observations up to time t

$$p(\mathbf{X}_t | \mathbf{Z}_t) \sim p(\mathbf{z}_t | \mathbf{x}_t, \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{x}_t | \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{X}_{t-1} | \mathbf{Z}_{t-1})$$

We assume the observations $\mathbf{z}_0, \mathbf{z}_1, \dots, \mathbf{z}_t$ are independent both

(1) mutually and (2) with respect to the dynamical process:

$$p(\mathbf{z}_t | \mathbf{x}_t, \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) = p(\mathbf{z}_t | \mathbf{x}_t)$$

We now have:

$$p(\mathbf{X}_t | \mathbf{Z}_t) \sim p(\mathbf{z}_t | \mathbf{x}_t) p(\mathbf{x}_t | \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{X}_{t-1} | \mathbf{Z}_{t-1})$$

Tracking over Time

\mathbf{x}_t : State at time t

$\mathbf{X}_t = \mathbf{x}_t, \mathbf{x}_{t-1}, \dots, \mathbf{x}_0$: States from time 0 to time t

\mathbf{z}_t : Observations performed at time t .

$\mathbf{Z}_t = \mathbf{z}_t, \mathbf{z}_{t-1}, \dots, \mathbf{z}_0$: Observations up to time t

$$p(\mathbf{X}_t | \mathbf{Z}_t) \sim p(\mathbf{z}_t | \mathbf{x}_t) p(\mathbf{x}_t | \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{X}_{t-1} | \mathbf{Z}_{t-1})$$

We want to get to:

$$p(\mathbf{x}_t | \mathbf{Z}_t) \sim p(\mathbf{z}_t | \mathbf{x}_t) \int_{\mathbf{x}_{t-1}} p(\mathbf{x}_t | \mathbf{x}_{t-1}) p(\mathbf{x}_{t-1} | \mathbf{Z}_{t-1})$$

Tracking over Time

\mathbf{x}_t : State at time t

$\mathbf{X}_t = \mathbf{x}_t, \mathbf{x}_{t-1}, \dots, \mathbf{x}_0$: States from time 0 to time t

\mathbf{z}_t : Observations performed at time t .

$\mathbf{Z}_t = \mathbf{z}_t, \mathbf{z}_{t-1}, \dots, \mathbf{z}_0$: Observations up to time t

$$p(\mathbf{X}_t | \mathbf{Z}_t) \sim p(\mathbf{z}_t | \mathbf{x}_t) p(\mathbf{x}_t | \mathbf{X}_{t-1}, \mathbf{Z}_{t-1}) p(\mathbf{X}_{t-1} | \mathbf{Z}_{t-1})$$

We can make \mathbf{x}_{t-1} appear:

$$p(\mathbf{X}_t | \mathbf{Z}_t) \sim p(\mathbf{z}_t | \mathbf{x}_t) p(\mathbf{x}_t | \mathbf{x}_{t-1}, \mathbf{X}_{t-2}, \mathbf{Z}_{t-1}) p(\mathbf{X}_{t-1} | \mathbf{Z}_{t-1})$$

We assume the states form a Markov chain: $p(\mathbf{x}_t | \mathbf{x}_{t-1}, \mathbf{X}_{t-2}, \mathbf{Z}_{t-1}) = p(\mathbf{x}_t | \mathbf{x}_{t-1})$

So:

$$p(\mathbf{X}_t | \mathbf{Z}_t) \sim p(\mathbf{z}_t | \mathbf{x}_t) p(\mathbf{x}_t | \mathbf{x}_{t-1}) p(\mathbf{X}_{t-1} | \mathbf{Z}_{t-1})$$



Tracking over Time

\mathbf{x}_t : State at time t

$\mathbf{X}_t = \mathbf{x}_t, \mathbf{x}_{t-1}, \dots, \mathbf{x}_0$: States from time 0 to time t

\mathbf{z}_t : Observations performed at time t .

$\mathbf{Z}_t = \mathbf{z}_t, \mathbf{z}_{t-1}, \dots, \mathbf{z}_0$: Observations up to time t

We had:

$$p(\mathbf{X}_t | \mathbf{Z}_t) \sim p(\mathbf{z}_t | \mathbf{x}_t) p(\mathbf{x}_t | \mathbf{x}_{t-1}) p(\mathbf{X}_{t-1} | \mathbf{Z}_{t-1})$$

Making \mathbf{X}_{t-2} appear:

$$p(\mathbf{x}_t, \mathbf{x}_{t-1}, \mathbf{X}_{t-2} | \mathbf{Z}_t) \sim p(\mathbf{z}_t | \mathbf{x}_t) p(\mathbf{x}_t | \mathbf{x}_{t-1}) p(\mathbf{x}_{t-1}, \mathbf{X}_{t-2} | \mathbf{Z}_{t-1})$$

$$\int_B p(A, B) = p(A)$$

Integrating over \mathbf{X}_{t-2} :

$$p(\mathbf{x}_t, \mathbf{x}_{t-1} | \mathbf{Z}_t) \sim p(\mathbf{z}_t | \mathbf{x}_t) p(\mathbf{x}_t | \mathbf{x}_{t-1}) p(\mathbf{x}_{t-1} | \mathbf{Z}_{t-1})$$



Tracking over Time

\mathbf{x}_t : State at time t

$\mathbf{X}_t = \mathbf{x}_t, \mathbf{x}_{t-1}, \dots, \mathbf{x}_0$: States from time 0 to time t

\mathbf{z}_t : Observations performed at time t .

$\mathbf{Z}_t = \mathbf{z}_t, \mathbf{z}_{t-1}, \dots, \mathbf{z}_0$: Observations up to time t


$$p(\mathbf{x}_t, \mathbf{x}_{t-1} \mid \mathbf{Z}_t) \sim p(\mathbf{z}_t \mid \mathbf{x}_t) p(\mathbf{x}_t \mid \mathbf{x}_{t-1}) p(\mathbf{x}_{t-1} \mid \mathbf{Z}_{t-1})$$

Integrating over \mathbf{x}_{t-1} :

$$p(\mathbf{x}_t \mid \mathbf{Z}_t) \sim p(\mathbf{z}_t \mid \mathbf{x}_t) \int_{\mathbf{x}_{t-1}} p(\mathbf{x}_t \mid \mathbf{x}_{t-1}) p(\mathbf{x}_{t-1} \mid \mathbf{Z}_{t-1})$$

Kalman Filter

$$p(\mathbf{x}_t \mid \mathbf{Z}_t) = N(\mathbf{x}_t; \mu_t, \Sigma_t)$$



Time update or Prediction

The states \mathbf{x}_t are assumed to evolve according to a dynamics model of the form:

$$\mathbf{x}_t = \mathbf{A} \mathbf{x}_{t-1} + \mathbf{w}_t$$

where:

- matrix \mathbf{A} is a state transition matrix;
- vector \mathbf{w}_t is the process noise.

State transition matrix A

Example:

If we assume the motion model is a constant velocity model, with no rotation:

$$\mathbf{x}_t = \begin{bmatrix} \mathbf{T}_t \\ \dot{\mathbf{T}}_t \end{bmatrix} \quad \mathbf{A} = \begin{bmatrix} \mathbf{I} & \mathbf{I} \\ \mathbf{0} & \mathbf{I} \end{bmatrix}$$

$$\begin{bmatrix} \mathbf{T}_{t+1} \\ \dot{\mathbf{T}}_{t+1} \end{bmatrix} = \begin{bmatrix} \mathbf{I} & \mathbf{I} \\ \mathbf{0} & \mathbf{I} \end{bmatrix} \begin{bmatrix} \mathbf{T}_t \\ \dot{\mathbf{T}}_t \end{bmatrix} + \mathbf{w}_t$$

The measurement update or Correction


Linear case:

The measurements \mathbf{z}_t are assumed to be related to the state \mathbf{x}_t by a linear measurement model:

$$\mathbf{z}_t = \mathbf{C} \mathbf{x}_t + \mathbf{v}_t$$

where

the vector \mathbf{v}_t represents the measurement noise.



This is not linear but it is always possible to linearize it.

Kalman filter

$$p(\mathbf{x}_t | \mathbf{Z}_t) \sim p(\mathbf{z}_t | \mathbf{x}_t) \int_{\mathbf{x}_{t-1}} p(\mathbf{x}_t | \mathbf{x}_{t-1}) p(\mathbf{x}_{t-1} | \mathbf{Z}_{t-1})$$


Under the assumptions that: $\mathbf{z}_t = \mathbf{C} \mathbf{x}_t + \mathbf{v}_t$ and $\mathbf{x}_t = \mathbf{A} \mathbf{x}_{t-1} + \mathbf{w}_t$

If we know $p(\mathbf{x}_{t-1} | \mathbf{Z}_{t-1}) = N(\mathbf{T}_{t-1}; \boldsymbol{\mu}_{t-1}, \boldsymbol{\Sigma}_{t-1})$, can we compute $p(\mathbf{x}_t | \mathbf{Z}_t)$?

The Kalman filter

1. Time update or Prediction
→ Uses a motion model

$$\begin{cases} \mu_t^- = \mathbf{A}\mu_{t-1} \\ \Sigma_t^- = \mathbf{A}\Sigma_{t-1}\mathbf{A}^\top + \Sigma_w \end{cases}$$




2. Measurement update or Correction

$$\begin{cases} \mu_t = \mu_t^- + \mathbf{G}_t(\mathbf{z}_t - \mathbf{C}\mu_t^-) \\ \Sigma_t = \Sigma_t^- - \mathbf{G}_t\mathbf{C}\Sigma_t^- \end{cases}$$


where \mathbf{G}_t is the Kalman gain computed as:

$$\mathbf{G}_t = \Sigma_t^- \mathbf{C}^\top (\mathbf{C}\Sigma_t^- \mathbf{C} + \Sigma_v)^{-1}$$

Kalman filter




Kalman filter



Particle Filter


$$p(\mathbf{x}_t \mid \mathbf{Z}_t) \equiv \{(\mu_{t,i}, w_{t,i})\}_i$$



Particle Filter




Particle filters




Time $t-1$


Particle filters




Particle filters



Particle filters



Particle filters



Efficient resampling of particles

